

MERLIN observations of GRS 1915+105 :

a progress report

R. P. Fender^a, S. T. Garrington^b, D. J. McKay^{b,c}, T. W. B. Muxlow^b,
G. G. Pooley^d, R. E. Spencer^b, A. M. Stirling^b, E. B. Waltman^e

^a*University of Amsterdam, The Netherlands*

^b*University of Manchester, Jodrell Bank, UK*

^c*Joint Institute for VLBI in Europe, The Netherlands*

^d*Mullard Radio Astronomy Laboratory, UK*

^e*Remote Sensing Division, NRL, USA*

Abstract

We present a progress report on MERLIN radio imaging of a radio outburst from GRS 1915+105. The major ejection occurred at the end of a ~ 20 day ‘plateau’ state, characterised by low/hard X-ray fluxes and a relatively strong flat-spectrum radio component. Apparent superluminal motions have been mapped with unprecedented resolution, and imply higher velocities in the jet than previously derived.

1. Introduction and Observations

We have observed the superluminal X-ray transient (Mirabel & Rodriguez 1994, hereafter MR94) with the MERLIN array in 1997 October – November following a major radio outburst.

The MERLIN observations were made with an angular resolution of 40 mas, five times better than previous VLA mapping, at a frequency of 5 GHz, on ten occasions over a period of 12 days. The radio flux from GRS 1915+105 was also simultaneously monitored at 2 & 8 GHz with the Green Bank Interferometer (GBI) and at 15 GHz with the Ryle Telescope (RT). Over the same period the system was monitored in soft X-rays with the Rossi XTE satellite.

2. X-ray and radio state

Prior to our MERLIN observations GRS 1915+105 had clearly been in an unusual state in both X-rays and radio, summarised in Fig. 1. Approximately 25 days prior to our first epoch of mapping, the source had undergone a significant radio flare, and then entered a ‘plateau’ state. This state is characterised by relatively bright and stable radio emission with a flat / inverted spectrum, and low but persistent X-ray emission with a hard spectrum. Previous monitoring reported at 15 GHz with the RT in Pooley & Fender (1997) and at lower frequencies in Bandyopadhyay et al. (1998) confirm the association between the plateau state and relatively bright radio, and perhaps infrared, emission.

3. Superluminal ejections

As indicated in Fig. 1, a sequence of ten MERLIN high-resolution radio maps were made of GRS 1915+105. The first four of these are presented in Fig. 2, revealing the clear expansion of the source on




Fig. 1. Radio (GBI) and X-ray (XTE) monitoring around the time of the MERLIN observations of GRS 1915+105. Around MJD 50725 a significant radio flare signals the beginning of the plateau state, characterised by relatively bright and stable flat-spectrum radio emission and steady, hard-spectrum emission in X-rays. The flare at the end of the plateau period triggered our MERLIN observations, indicated by diamonds on the top panel. Images from the first four epochs of MERLIN observations are presented in Fig. 2.




Fig. 2. Radio maps of GRS 1915+105 from the first four epochs of our MERLIN observations. The observing frequency is 5 GHz (6 cm) and the angular resolution 40 mas. In the first two epochs the core and receding component are blended together; by the third epoch core and approaching (SE) and receding (NW) components are clearly resolved. The proper motions are significantly higher than those reported in MR94 (see text). The unit contour level for each epoch is indicated on the right side of the figure, and the maps have been rotated clockwise by ~ 145 degrees for clarity.

daily timescales. The jet sidedness is consistent with that reported in MR94, and we are able to track the proper motions of three approaching and one receding components over the entire set of images. The position angle of the ejections, approximately 145 degrees, is also very similar to that reported in MR94.

All ejections are consistent with ballistic motions to better than 10%, and we find best-fits to the proper motions of :

$$\mu_{\text{app}} = 23.6 \pm 0.5$$

and

$$\mu_{\text{rec}} = 10.0 \pm 0.5$$

mas. d^{-1} , significantly greater than those reported in MR94. All fits are good, with $\chi^2_{\text{red}} \leq 1$. The proper motion of $17.6 \pm 0.4 \text{ mas. d}^{-1}$ reported by MR94 for the approaching component can be ruled out for this ejection; fixing the proper motion to this value does not give an acceptable fit to the data.

Initial analysis of the proper motions, under the standard assumption of an intrinsically symmetric ejection, suggests that we are measuring significantly greater velocities (by $\geq 10\%$) than MR94, at a similar angle to the line of sight (most likely in the range 60 – 70 degrees).

Again assuming a symmetric ejection, we can use the proper motions to place limits on the distance to GRS 1915+105. At the most extreme, given the measurement uncertainties, the distance must be ≤ 13.6 kpc. A more likely upper limit is ≤ 11.2 kpc, i.e. significantly closer (although within their estimated errors) than the best estimate of 12.5 kpc used by MR94.

4. Conclusions

We have presented preliminary results from a major set of MERLIN observations of GRS 1915+105 simultaneous with multiwavelength radio and X-ray monitoring. While the nature of the plateau state itself remains unclear, it is now certain that the radio flaring associated with the end of the state **is** associated with relativistic ejections.

These ejections, in particular the approaching component, have significantly higher proper motions than those recorded by MR94, most simply interpreted as higher intrinsic velocities at a similar angle to the line of sight. We also find that GRS 1915+105 is likely to be significantly closer than the best estimate of 12.5 kpc of MR94.

More details of the observations, including spatially resolved linear polarisation maps, and their interpretation, will be found in a paper to be submitted to MNRAS.

Acknowledgements

We acknowledge great efforts by many people, mostly at Jodrell Bank, which helped to make these observations a success. In particular we would like to thank Peter Thomasson, Shane McKie and Richard Ogley. MERLIN is operated by the University of Manchester at the Nuffield Radio Astronomy Laboratories, Jodrell Bank, on behalf of the Particle Physics and Astronomy Research Council. We acknowledge also the use of public data from XTE ASM and GBI. RPF acknowledges support from EC Marie Curie Fellowship ERBFMBICT 972436, and DJM from EC TMR-LSF contract No. ERBFMGECT 950012.

References

- Bandyopadhyay R., Martini P., Gerard E., Charles P.A., Wagner R.M., Shrader C., Shahbaz T., Mirabel I.F., 1998, MNRAS, 295, 623
- Mirabel I.F., Rodriguez L.F., 1994, Nature, 371, 46
- Pooley G.G., Fender R.P., 1997, MNRAS, 292, 925